

Dragonflies from mainland Yemen and the Socotra Archipelago – additional records and novelties

Wolfgang Schneider¹ and Abdul Karim Nasher²

¹Entomology II, Research Institute and Natural History Museum Senckenberg, Senckenberganlage 25, 60325 Frankfurt a.M., Germany. Email: fri.wol@t-online.de
wolfgang.schneider@senckenberg.de

²Faculty of Science, University of Sana'a, University Post Office, P.O. Box 12231, Sana'a, Yemen. Email: karimnasher@yahoo.com

Abstract

The odonatological results of two field trips to mainland Yemen carried out mainly in summer 2005 and winter 2007, and to Socotra in winter 1999 are put on record. At 30 localities, 33 dragonfly species were collected, respectively observed (3). One species, *Azuragrion somalicum* is new for mainland Yemen, and a second, *Pseudagrion niloticum*, is new for mainland Yemen and the Arabian Peninsula. Three species, *Azuragrion somalicum*, *Orthetrum julia*, and *Sympetrum fonscolombii* are new records for Socotra.

Kurzfassung

Die odonatologischen Ergebnisse zweier Feldaufenthalte auf dem jemenitischen Festland hauptsächlich im Sommer 2005 und Winter 2007 und auf Socotra im Winter 1999 werden mitgeteilt. An 30 Fundorten wurden 33 Libellenarten gesammelt oder beobachtet (3). *Azuragrion somalicum* ist neu für das jemenitische Festland, *Pseudagrion niloticum*, ist neu für das jemenitische Festland und die Arabische Halbinsel. *Azuragrion somalicum*, *Orthetrum julia* und *Sympetrum fonscolombii* sind Neunachweise für Socotra.

Introduction

The southwest of Arabia, comprising the Asir Mountains in Saudi Arabia and the high mountains of Yemen, is one of three biological hotspots of the peninsula, the other two being Dhofar (Oman) and Hadhramout (Yemen). These are considered relict pockets from more humid spells of the late Pleistocene and Holocene (Samraoui et al. 1993). With more than 40 taxa the southwest exhibits the highest species richness.



Although the three regions seem to differ slightly in species composition, this may turn out as an artifact once the odonate fauna of the Mahri and al-Hawf regions close to Dhofar become better known. Compared to other oceanic islands and taking into account its large size, Socotra has an impoverished odonate fauna with only one endemic.

Abbreviations

Collectors and/or illustrators: E.F.: Eva Feldkamp (Frankfurt a.M., Germany). – A.K.N.: Abdul Karim Nasher (University of Sana'a, Yemen). – H.P.: Hans Pohl (Phyletisches Museum, University of Jena, Germany). – W.S.: Wolfgang Schneider (Senckenberg Museum, Frankfurt a.M., Germany). – K.v.D.: Kay van Damme (University of Birmingham, UK).

List of localities with species present

Yemen mainland (localities 1-16) (Figure 1)

1. Wadi al-Khun (Hadhramout): 03.06.2005, 16°09'45"N 49°04'46"E, 689 m a.s.l.: *Arabicnemis caerulea*, *Ischnura evansi*, *Trithemis annulata*, *T. arteriosa*




Fig. 1: Map of mainland Yemen, showing position of collecting sites (E. F.).



2. Wadi al-Khun (Hadhramout): 03.06.2005, 16°09'45"N 49°06'02"E, 659 m a.s.l.: *Arabicnemis caerulea*, *Ceriagrion glabrum*, *Ischnura evansi*, *Pseudagrion hamoni*, *Nesciothemis farinosa*, *Orthetrum abbotti*, *O. chrysostigma*, *Trithemis annulata*, *T. arteriosa*
3. Wadi Massila at Bir (Hadhramout): 04.06.2005, 16°08'36"N 49°27'07"E, 542 m a.s.l.: *Arabicnemis caerulea*, *Ischnura evansi*, *Crocothemis erythraea*, *Nesciothemis farinosa*, *Trithemis annulata*
4. Wadi Massila at al-Hind (Hadhramout): 05.06.2005, 15°44'53"N 50°24'32"E, 388 m a.s.l.: *Arabicnemis caerulea*, *Nesciothemis farinosa*, *Orthetrum sabina*, *Trithemis annulata*, *T. arteriosa*
5. Wadi Massila (Hadhramout): 05.06.2005, 15°43'22"N 50°27'23"E, 383 m a.s.l.: *Arabicnemis caerulea*, *Pseudagrion niloticum*, *Nesciothemis farinosa*
6. Wadi Ghail Omar in Wadi Sah system (Hadhramout): 02.06.2005, 15°40'51"N 48°51'59"E, 725 m a.s.l.: *Arabicnemis caerulea*, *Ischnura evansi*, *Pseudagrion hamoni*, *P. niloticum*, *P. sublacteum*, *Nesciothemis farinosa*, *Orthetrum chrysostigma*, *Trithemis annulata*, *T. kirbyi*
7. Wadi Adhm/Ghail Omar in Wadi Sah system (Hadhramout): 02.06.2005, 15°40'10"N 48°51'04"E, 727 m a.s.l.: *Arabicnemis caerulea*, *Ceriagrion glabrum*, *Ischnura evansi*, *Pseudagrion hamoni*, *P. niloticum*, *Nesciothemis farinosa*
8. Al-Ahjar village: artificial pond fed by waterfall, 10.06.2005 & 25.10.2007, 15°28'19"N 43°51'26"E, 2487 m a.s.l.: *Pseudagrion arabicum*, *Pinheyschna yemenensis*, *Crocothemis sanguinolenta*, *Orthetrum caffrum*, *O. julia*
9. Al-Ahjar: artificial pond fed by spring water, 10.06.2005 & 25.10.2007, 15°27'49"N 43°53'36"E, 2407 m a.s.l.: *Anax imperator*, *Crocothemis erythraea*, *Orthetrum caffrum*, *Sympetrum fonscolombii*
10. Matnah, 30 km W of Sana'a: 30.01.1999, 15°14'N 43°60'E, 2700 m a.s.l.: *Sympetrum fonscolombii*
11. Bani Matar: 06.03.2008, 15°12'07"N 43°59'34"E, 2720 m a.s.l.: *Orthetrum caffrum*
12. Wadi Mara in Wadi Dau'an system (Hadhramout): 31.05.2005, 15°08'36"N 48°26'55"E, 1002 m a.s.l.: *Arabicnemis caerulea*, *Pseudagrion hamoni*, *Nesciothemis farinosa*, *Trithemis dejouxi*, *T. kirbyi*
13. Manaka (W of): 21.12.2007, 15°07'17"N 43°37'42"E, 962 m a.s.l.: *Zygonyx torridus*
14. Ad-Dawasir (Hadhramout): 30.05.2005, stagnant pool, 14°56'03"N 48°55'07"E, 1264 m a.s.l.: *Azuragrion somalicum*, *Ischnura evansi*, *Tamea limbata*, *Trithemis arteriosa*
15. Ghail Bawazir (Hadhramout): 07.12.2007, 14°47'31"N 49°22'46"E, 124 m a.s.l.: *Ischnura evansi*, *Pseudagrion hamoni*



16. Abyan, Batais village, 08.11.2007, 13°21'00"N 45°17'46"E, 170 m a.s.l.: *Anax imperator*, *Hemianax ephippiger*, *Paragomphus genei*, *Crocothemis erythraea*, *Pantala flavescens*, *Trithemis annulata*, *T. arteriosa*, *T. kirbyi*

Socotra island and Samha (localities 17-30) (Figure 2)

17. Hadibo: town and surroundings, 02.02.-26.02.1999, 12°39'11"N 54°01'08"E, 7 m a.s.l.: *Macrodiplax cora*
18. Wadi Daneghan: 13.02.1999, 12°38'59"N 54°01'04"E, 10 m a.s.l.: *Trithemis arteriosa*




Fig. 2: Map of the Socotra Archipelago, showing position of collecting sites on Socotra and Samha (E.F.).

19. Coastal road in eastern direction: 04.02.1999, 12°37'56"N 54°09'16"E, 31 m a.s.l.: *Ceriagrion glabrum*, *Ischnura senegalensis*, *Zygonyx torridus*
20. Wadi Zeewef, ascent to Homhil: ponds between shrubs, 07.02.1999, 12°35'36"N 54°18'53"E, 136 m a.s.l.: *Orthetrum chrysostigma*
21. Wadi Zeewef, Homhil plain: 07.-08.1999, 12°35'16"N 54°18'09"E, 337 m a.s.l.: *Azuragrion granti*, *A. nigradorsum*, *Ceriagrion glabrum*, *Anax imperator*, *Crocothemis erythraea*, *Diplacodes lefebvrii*, *Orthetrum chrysostigma*, *Trithemis arteriosa*
22. Wadi Daneghan, ascent to Adho Dimelho: 02.02.2005, 12°34'N 54°02'E, 1190 m a.s.l.: *Azuragrion granti*, *Crocothemis erythraea*, *Trithemis arteriosa*
23. Adho Dimelho: small river between shrubs, 03.02.1999, 12°34'34"N 54°02'55"E, 940 m a.s.l.: *Azuragrion granti*, *Orthetrum julia*, *Trithemis arteriosa*
24. Wadi in the Haggier Mountains: 24.02.1999, 12°33'31"N 54°01'E, 1347 m a.s.l.: *Azuragrion granti*, *Anax imperator*, *Orthetrum chrysostigma*, *Trithemis arteriosa*
25. Deksam plateau: 22.-24.02.1999, 12°32'26"N 53°59'21"E, 1020 m a.s.l.: *Azuragrion granti*, *A. nigradorsum*, *Trithemis arteriosa*



26. Deksam plateau: shallow marsh, 22.-24.02.1999, 12°32'26"N 53°59'43"E, 1010 m a.s.l.: *Azuragrion somalicum*
27. Deksam plateau: 22.02.1999, 12°31'26"N 53°57'11"E, 1000 m a.s.l.: *Orthetrum chrysostigma*, *Trithemis arteriosa*
28. Wadi Kilisan canyon: 10.02.1999, 12°29'12"N 54°19'40"E, 150 m a.s.l.: *Azuragrion nigradorsum*, *Crocothemis erythraea*, *Diplacodes lefebvreii*, *Sympetrum fonscolombii*, *Trithemis arteriosa*
29. Wadi en route to Nogad: 14.02.1999, 12°26'27"N 54°09'19"E, 180 m a.s.l.: *Ischnura senegalensis*, *Crocothemis erythraea*, *Trithemis arteriosa*
30. Samha Island: well, 16.02.1999, 12°10'N 53°01'E, 170 m a.s.l.: *Trithemis arteriosa*

Systematic account

Platycnemididae - Featherlegs

Arabicnemis caerulea Waterston, 1984 – Powder Blue Damsel

Specimens studied: Mainland: 1: 2 ♂♂, 1 ♀ (1 pair in copula), W.S. – 2: 11 ♂♂, 1 ♀♀, W.S. – 3: 3 ♂♂, 1 ♀, W.S. – 4: 4 ♂♂, 2 ♀♀, W.S. – 5: 3 ♂♂, W.S. – 6: 1 ♂, W.S. – 7: 2 ♂♂, 1 ♀ (1 pair in copula), W.S. – 12: 8 ♂♂, 3 ♀♀ (one pair in copula).

In 2005 *Arabicnemis caerulea* was abundant at all its collecting sites in Wadi Hadhramout. The most recent records (2008-2011) are localized in the northeastern subpopulation in Oman (Wilson 2008, Schneider unpublished).

Coenagrionidae – Pond Damsels

Azuragrion granti (McLachlan, 1903) – Grant's Bluet

Specimens studied: Socotra: 21: 1 ♂, 1 ♀, H.P. – 22: 2 ♂♂, 1 ♀, H.P. – 23: 1 ♀, H.P. – 24: 4 ♂♂, 2 ♀♀. – 25: 4 ♂♂, H.P.

Originally described in the genus *Ischnura* Charpentier, 1840, the species was later transferred to *Enallagma* Charpentier, 1840 by McLachlan (1903) and finally placed in a new genus *Azuragrion* by May (2002). This species, the only known endemic odonate of Socotra, seems to be restricted to the mountainous eastern part of the island (Riservato et al. 2010).

Azuragrion nigradorsum (Selys, 1876) – Sailing Bluet

Specimens studied: Socotra: 21: 5 ♂♂, 1 ♀, H.P. – 28: 1 ♂, H.P.

In Arabia the species has a discontinuous range with two areas in mainland Yemen and southern Oman and one in the eastern part of Socotra island.

Azuragrion somalicum (Longfield, 1931) [subsp. *amitinum* Waterston, 1991] – Somali Bluet

Specimens studied: 14: 1 ♂, W.S. – 26: 1 ♂, K.v.D.



New to mainland Yemen and the Socotra Archipelago. Although the species' presence in Yemen was anticipated (Schneider & Dumont 1997), this record lies about 600 km west of its known distribution around Salalah in Dhofar, Oman. The most recent Arabian record published together with a good photograph of a male is from a Wadi NE of Salalah (van der Weide & Kalkman 2008). The new record from Hadhramout suggests a wider distribution in south Arabia. To clarify the status of the subspecies *amitinum* Waterston, 1991 longer series of specimens are needed.

Ceriagrion glabrum (Burmeister, 1839) – Common Citril

Specimens studied: Mainland: 2: 5 ♂♂, 1 ♀, W.S. – 7: 1 ♂, W.S. – Socotra: 19: 6 destroyed specimens, H.P. – 21: 2 ♂♂, H.P.

Ischnura evansi Morton, 1919 – Evan's Bluetail

Specimens studied: Mainland: 1: 1 ♀, W.S. – 2: 1 ♀, W.S. – 3: 3 ♂♂, 2 ♀♀, W.S. – 6: 1 ♂, 1 ♀, W.S. – 7: 1 ♂, W.S. – 14: 4 ♂♂, 1 ♀, A.K.N. – 15: 3 ♂♂, 1 ♀, A.K.N.

Ischnura senegalensis (Rambur, 1842) – Marsh Bluetail

Specimens studied: Socotra: 19: 1 ♂, H.P. – 29: 1 ♂, H.P.

Pseudagrion arabicum Waterston, 1980 – Arabian Sprite

Specimens studied: Mainland: 8: 1 ♂ (2007), W.S. & A.K.N.

Pseudagrion arabicum is a rare endemic species of southwestern Arabia and is known from a rather small range in the coastal mountains of the Red Sea in Saudi Arabia and Yemen. This is a high mountain species ranging from 2,000 to 3,000 m a.s.l. Ten records are now available from 6 different localities (Waterston 1980, 1985; Dumont & Al-Safadi 1991; Schneider & Krupp 1993; Schneider & Parr 1998).

Pseudagrion hamoni Fraser, 1955 – Hamon's Sprite

Specimens studied: Mainland: 2: 2 ♂♂, W.S. – 6: 17 ♂♂, 3 ♀♀, W.S. – 7: 5 ♂, 1 ♀, W.S. – 12: 7 ♂♂, 3 ♀♀, W.S. – 15: 1 ♂, A.K.N.

Pseudagrion hamoni is widely distributed in sub-Saharan Africa, Somalia, and Ethiopia, with Sahelian relict populations in Mauritania, S-Algeria, S-Libya. In Arabia the species is common in the mountains of the south and southwest of Yemen and Saudi Arabia. 70 records are now available from 45 different localities. (Waterston 1985; Al-Safadi 1990; Dumont & Al-Safadi 1991, 1993; Schneider & Krupp 1993; Krupp et al. 2006; Lambret & Boudot 2009).

Pseudagrion niloticum Dumont, 1978 (Figure 3) – Nile Sprite

Specimens studied: Mainland: 5: 11 ♂♂, 2 ♀♀, W.S. – 6: 8 ♂♂, 1 ♀, W.S. – 7: 5 ♂♂, 1 ♀, W.S. Comparative specimens: males from Kenya (Coll. V. Clausnitzer & Coll. K.-D.B. Dijkstra), males and a female from Sudan (Coll. H.J. Dumont)



This species is new for Yemen and the Arabian Peninsula and an interesting addition from the zoogeographical point of view. It is considered the northern vicariant (Nile valley, Somalia, East Ethiopia) of *P. acacia* (southern Africa, African Plateau) with which it was often confused in the past (Dumont 1978). In fact, both species are so closely related that Dumont & Martens (1984) postulated a common, late Pleistocene ancestor.




Fig. 3: *Pseudagrion niloticum*, male from Wadi Massila (collecting station 5). a: tip of abdomen with anal appendages in lateral view. b: the same in dorsal view (drawings W.S.).

Pseudagrion sublacteum (Karsch, 1893) – Cherry-eye Sprite

Specimens studied: Mainland: 6: 4 ♂♂, W.S.

In Arabia *Pseudagrion sublacteum* is restricted to Yemen, southern Oman and the western Red Sea mountains in Saudi Arabia (Waterston 1980, Waterston & Pittaway 1991, Schneider & Krupp 1993) with voucher specimen checked and confirmed by the first author. 42 records are available from 22 localities. Arabian populations were first ascribed to the Levantine subspecies *mortoni* Schmidt, 1936, but a comparison with African populations has shown that Arabian populations belong to the nominal subspecies (Dumont & Al-Safadi 1991).



Aeshnidae - Hawkers

Anax imperator (Leach, 1815) – Blue Emperor

Specimens studied/observation: Mainland: 9: 1 ♂ (observation, 2005), W.S. – 16: 1 ♂, 2 ♀, A.K.N. – Socotra: 21: 1 ♀, caught in Malaise trap at night, H.P. – 24: 1 ♀, H.P.

Hemianax ephippiger (Burmeister, 1839) – Vagrant Emperor

Specimens studied: Mainland: 16: 2 ♂♂, A.K.N.

Pinheyschna yemenensis (Waterston, 1985) – Arabian Hawker

Observations: Mainland: 8: Several specimens seen hovering high in the air (2007).

Gomphidae – Clubtails

Paragomphus genei (Selys, 1841) – Green Clubtail

Specimens studied: Mainland: 16: 2 ♂♂, 3 ♀♀, A.K.N.

Libellulidae - Skimmers

Crocothemis erythraea (Brullé, 1832) – Broad Scarlet

Specimens studied: Mainland: 3: 1 male, W.S. – 9: 1 male, 2 females (2005), W.S. – 16: 1 male, 1 female, A.K.N. – Socotra: 21: 2 ♂♂, H.P. – 22: 1 ♂, H.P. – 28: 2 ♂♂, H.P. – 29: 1 ♂, H.P.

Crocothemis sanguinolenta (Burmeister, 1839) – Little Scarlet

Specimens recorded: Mainland: 8: 1 ♂ (2005), W.S.

Diplacodes lefebvrii (Rambur, 1842) – Black Percher

Specimens studied: Socotra: 21: 1 ♂, 1 ♀, H.P. – 28: 2 ♂♂, H.P.

Nesciothemis farinosa (Förster, 1891) – Black-tailed Skimmer

Specimens studied: Mainland: 2: 1 ♂, W.S. – 3: 1 ♂, W.S. – 4: 1 ♂, W.S. – 5: 1 ♂, W.S. – 6: 6 ♂♂, W.S. – 7: 1 ♀, W.S. – 12: 1 ♂, W.S.

Macrodiplax cora (Kaup in Brauer, 1867) – Coastal Glider

Specimens studied: Socotra: 17: 1 ♂, H.P.

Orthetrum abbotti (Calvert, 1892) – Abbott's Skimmer

Specimens studied: Mainland: 2: 3 ♂♂, W.S.

This new locality and the first one, Hami hotspots (Dumont & al-Safadi 1993), lie in Hadhramout. As the species has been recorded from Jordan (Dumont 1977, Monnerat 2010), it should be searched for in wadi systems along the Red Sea coast of Saudi Arabia and Yemen.

Orthetrum caffrum (Burmeister, 1839) – Two-striped Skimmer

Specimens studied: Mainland: 8: 7 ♂♂ (2005), W.S., 4 ♂♂ (2007), W.S. & A.K.N. – 9: 1 ♀ (2005), W.S., 1 ♂ (2007), W.S. & A.K.N. – 11: 1 ♂, A.K.N.



Orthetrum chrysostigma (Burmeister, 1839) – Epaulet Skimmer

Specimens studied: Mainland: 2: 1 ♂, W.S. – 6: 1 ♂, W.S. – Socotra: 20: 1 ♂. – 21: 2 ♂♂, H.P. – 24: 1 ♂, 1 ♀, H.P. – 27: 2 ♂♂, 1 ♀, H.P.

Orthetrum julia Kirby, 1900 [*falsum* Longfield, 1955] – Julia Skimmer

Specimens studied: Mainland: 8: 2 ♂♂ (2007), W.S. & A.K.N. – Socotra: 23: 1 ♂, H.P.

This is a new record for Socotra. On the mainland, the species is known only from one locality in the west of Yemen, al-Ahjar (collecting station 8) where it was discovered in March 1996 (Schneider and Parr 1998).

Orthetrum sabina (Drury, 1773) – Slender Skimmer

Observation: Mainland: 4: 1 ♂ observed, W.S.

Pantala flavescens (Fabricius, 1798) – Globe Skimmer

Specimens studied: Mainland: 16: 1 ♂, A.K.N.

Sympetrum fonscolombii (Selys, 1840) – Red-veined Darter

Specimens studied: Mainland: 9: 4 ♂♂ (2007), W.S. & A.K.N. – 10: 1 ♂, H.P. – 28: 2 ♂, H.P.

New record for Socotra (see Discussion).

Tramea limbata (Desjardins, 1832) – Voyaging Glider

Specimens studied: Mainland: 14: 1 ♂, W.S.

Trithemis annulata (Beauvais, 1807) – Violet Dropwing

Specimens studied: Mainland: 1: 1 ♂, W.S. – 2: 1 ♂, 1 ♀, W.S. – 3: 13 ♂♂, 2 ♀♀, W.S. – 4: 2 ♂♂, W.S. – 6: 3 ♂♂, W.S. – 16: 5 ♂♂, A.K.N.

Trithemis arteriosa (Burmeister, 1839) – Red-veined Dropwing

Specimens studied: Mainland: 1: 1 ♂ male, W.S. – 2: 6 ♂♂, 4 ♀, W.S. – 14: 3 ♂♂, W.S. – 16: 4 ♂♂, A.K.S. – Socotra: 18: 1 ♂, H.P. – 21: 1 ♂, H.P. – 22: 3 ♀♀, H.P. – 23: 1 ♂, H.P. – 24: 1 ♂, H.P. – 25: 1 ♂, H.P. – 27: 1 ♂, H.P. – 28: 1 ♂, 1 ♀, H.P. – 29: 2 ♂♂, H.P. – 30: 2 ♂♂, H.P.

Trithemis dejouxi Pinhey, 1978 – Dejoux's Dropwing

Specimens studied: Mainland: 12: 3 ♂♂, W.S.

The species was first recorded for Arabia and Yemen sub *Trithemis donaldsoni* by Dumont and Al-Safadi (1991) and again by Krupp et al. (2006) as *Trithemis* aff. *donaldsoni*. Structural characters are indeed so minor that Pinhey (1978) described the taxon as a northeastern subspecies of the in other parts of Africa widely distributed *T. donaldsoni* (Calvert, 1899). O'Neil & Paulson (2001) finally found stable characters to raise *T. dejouxi* to specific rank; they were followed by Dijkstra (2007) and Damm et al. (2010).



Trithemis kirbyi Selys, 1891 – Kirby's Dropwing

Specimens studied: Mainland: 6: 2 ♂♂, W.S. – 12: 1 ♂, W.S. – 16: 1 ♂, A.K.N.

Zygonyx torridus (Kirby, 1889) – Ringed Cascader

Specimens studied: Mainland: 13: 1 ♂, 3 ♀♀, A.K.N. – Socotra: 19: 1 ♀, H.P.

Discussion

Despite political and tribal instability, and a resulting decrease in field activities, it is the southwest of Arabia accounting for all recent additions to the odonate fauna of the peninsula. The steady increase in numbers of species in the last two decades is remarkable, and this trend is likely to continue: 52 species in Waterston & Pittaway (1991), 56 in Schneider & Krupp (1993), 57 in Dumont & Al-Safadi (1993), 59 in Schneider & Dumont (1997), 60 in Schneider & Parr (1998), and finally 61, when Dumont & Verschuren (2005) concluded that African and SW-Arabian *Orthetrum* so far identified as *taeniolum* Schneider, 1845 pertain to a separate sister taxon, *kollmannspergeri* Buchholz, 1959. The present paper raises the number of Arabian species to 62, and species records from mainland Yemen from 40 to 42.

As far as Socotra is concerned, in the light of increasing collecting efforts, frequent and long term visits by odonatologists in the last decade, the number of new species records (3) is rather poor and not as high as anticipated. Although additions can never be ruled out, the present list is close to be complete. *Sympetrum fonscolombii* is a notorious migrant with a wide distribution in the paleotropics. Likewise *Orthetrum julia* is widespread in Africa and was recently also found in SW-Arabia (Schneider & Parr 1998), both species were therefore expected to occur on the island. The presence of *Azuragrion somalicum*, known from E-Africa (Ethiopia, Somalia) and from two relict pockets in Oman and Yemen, is more of a surprise. As *A. somalicum* and *A. nigradorsum* are both very small species, a wind-borne dispersal and arrival on the island is not unlikely. The somewhat atypical (size, morphology) *Azuragrion granti* is restricted to the eastern mountainous part of the island and may well represent a relict of an old fauna. Keeping in mind that absence of evidence is not evidence of absence, it is striking that the genus *Pseudagrion* Selys, 1876 is obviously not represented on Socotra, but is on other western Indian Ocean islands (whether continental or oceanic) like Madagascar, the Comores, and the Mascarenes. A major freshwater crisis on Socotra in the past with species extinctions is a possible explanation. The absence of primary freshwater fish and amphibians points in the same direction.

Acknowledgements

Special thanks go to Hans Pohl (University of Jena, Germany) and Kai van Damme (University of Birmingham, UK) for collecting Odonata on Socotra. Friedhelm Krupp (Qatar Museum Authority, Doha, Qatar), A. Miller (Royal Botanic Garden, Edinburgh, UK),



and Uwe Zajons (BiK-F & Senckenberg Research Institute, Frankfurt, Germany), and Eduardo Zandri (UNDP, Nairobi, Kenya) are acknowledged for their logistic support in the field. We are grateful to V. Clausnitzer (Halle, Germany), K.D.-B. Dijkstra (NMNH Naturalis, Leiden, The Netherlands), and H.J. Dumont (University of Gent, Belgium) for their loan of African specimens of *Pseudagrion niloticum* for comparison.

References

*Relevant literature not mentioned in the text are marked with an asterisk.

Al-Safadi, M.M. 1990. Dragonflies (Odonata) of the Yemen Arab Republic. Fauna of Saudi Arabia 11: 18-30.

*Al-Safadi, M.M. 1995. A pilot study of lake Ma'rib, Yemen. Hydrobiologia 315: 203-209.

*Bradshaw, C.G. 1996. Odonata observations in Yemen, spring 1993. Sandgrouse 17: 188.

Carfi, S. 1974. Contribution to the knowledge of Somalian Odonata. Monitore zoologico italiano (N.S.) Suppl. 5: 147-181.

*Carfi, S., R. Vittoria & F. Terzani 1995. Some dragonflies from the north of the Republic of Yemen. Bolletino della Società entomologica italiana 126: 195-199.

Clausnitzer, V. & K.-D.B. Dijkstra 2005. The dragonflies (Odonata) of Ethiopia, with notes on the status of endemic taxa and the description of a new species. Entomologische Zeitschrift 115: 117-130.

Damm, S., K.-D.B. Dijkstra & H. Hadrys 2010. Red drifters and dark residents: the phylogeny of a Plio-Pleistocene dragonfly radiation reflects Africa's changing environment (Odonata, Libellulidae, *Trithemis*). Molecular Phylogenetics and Evolution 54: 870-882.

Dijkstra, K.-D.B. 2007: The name-bearing types of Odonata held in the Natural History Museum of Zimbabwe, with systematic notes on Afrotropical taxa. Part 1: Introduction and Anisoptera. International Journal of Odonatology 10: 1-29.

Dumont, H.J. 1977. *Orthetrum abbotti* Calvert, 1892, a new Ethiopian representative in the Palaearctic fauna (Anisoptera: Libellulidae). Odonatologica 6: 199-203.

Dumont, H.J. 1978. On confusion about the identity of *Pseudagrion acacia* Foerster, 1906, with description of *P. niloticum* spec. nov., and on the identity of *P. hamoni* Fraser, 1955 (Zygoptera: Coenagrionidae). Odonatologica 7: 123-133.

Dumont, H.J. & M.M. Al-Safadi 1991. Additions to the dragonfly fauna of Yemen. Notulae odonatologicae 3: 114-117.

Dumont, H.J. & M.M. Al-Safadi 1993. Further additions to the dragonfly fauna of the Republic of Yemen (Odonata). Opuscula zoologica fluminensia 109: 1-8.



- Dumont, H.J. & K. Martens 1984. Dragonflies (Insecta: Odonata) from the Red Sea hills and the main Nile in Sudan. *Hydrobiologia* 110: 181-190.
- Dumont, H.J. & D. Verschuren 2005. Odonata from the Ennedi and Ounianga regions of northern Chad, with a note of the status of *Orthetrum kollmannspergeri* Buchholz, and a checklist of the species currently known from the Republic of Chad. *Odonatologica* 34(3): 291-297.
- Kimmins, D.F. 1960. The Odonata and Neuroptera of the Island of Socotra. *Annals and Magazine of Natural History, series 13*, 3(7): 385-392.
- Krupp, F., M. Apel, A. Hamoud, W. Schneider & U. Zajonz 2006. Zoological survey in the Red Sea coastal zone of Yemen. *Fauna of Arabia* 21: 11-32.
- Lambret, Ph J.-P. Boudot 2009. *Nesciothemis farinosa* (Förster, 1898) et *Orthetrum ransonnetii* (Brauer, 1865) nouveaux pour l'Arabie Saoudite et autres observations d'odonates sur les reliefs côtiers de la Mer Rouge. *Martinia* 25(4): 153-155.
- May, L.M. 2002. Phylogeny and taxonomy of the damselfly genus *Enallagma* and related Taxa (Odonata: Zygoptera: Coenagrionidae). *Systematic Entomology* 27: 387-408.
- McLachlan, R. 1903. The dragonflies of Sokotra. In: H.O. Forbes (ed.), *The Natural History of Sokotra and Abd-el-Kuri: being the report upon the results of the con joint expedition to these islands in 1898-9*. Special Bulletin of the Liverpool Museum, London, XLVII + 598 pp.
- Monnerat, C. 2010. *Crocothemis sanguinolenta* (Burmeister, 1839) and *Orthetrum abbotti* Calvert, 1890 in Jordan: distribution and ecological data. 29. Jahrestagung der GdO, Rothenburg ob der Tauber.
- O'Neil, G.O. & D.R. Paulson 2001. An annotated list of Odonata collected in Ghana in 1997, a checklist of Ghana Odonata, and comments on west African odonate biodiversity and biogeography. *Odonatologica* 30: 67-86.
- *Peters, G. & G. Theischinger 2011. The genera of the Afrotropical "Aeschnini": *Afro-aeschna* gen. nov., *Pinheyschna* gen. nov. and *Zosteraeschna* gen. nov. with the description of *Pinheyschna waterstoni* spec. nov. (Anisoptera: Aeschnidae). *Odonatologica* 40: 227-249.
- Pinhey, E. 1978. Comparative notes on African species of *Trithemis* Brauer (Odonata: Anisoptera) and its congeners. *Arnoldia Rhodesia* 8(26): 1-8.
- Riservato, E., C. Grieco, F. Pella, R. Sindaco, F. Pupin, A.S. Suleiman & M. Fasola 2010. A contribution to the knowledge of the odonatofauna of the Socotra Archipelago (Yemen). *Zoology in the Middle East* 50: 101-106.
- Samraoui, B., S. Benjacoub, S. Mecibah & H.J. Dumont 1993. Afrotropical libellulids in the lake district of El Kala, North-East Algeria, with a rediscovery of *Urothemis e. edwardsii* (Selys) and *Acisoma panorpoides ascalaphoides* (Rambur) (Anisoptera: Libellulidae). *Odonatologica* 22(3): 365-372.



- *Schneider, W. 1999. Soqatra – the island of the dragon’s blood. *Agrion* 3: 27-28.
- Schneider, W. & F. Krupp 1993. Dragonfly records from Saudi Arabia, with an annotated checklist of the species from the Arabian Peninsula (Insecta: Odonata). *Fauna of Saudi Arabia* 13: 63-78.
- Schneider, W. & H.J. Dumont 1997. The dragonflies and damselflies of Oman (Insecta: Odonata) of Oman. An updated and annotated checklist. *Fauna of Saudi Arabia* 16: 89-110.
- Schneider, W. & H.J. Dumont 1998. Checklist of the dragonflies and damselflies of Soqatra island (Insecta: Odonata). In: H.J. Dumont (ed.), *Proceedings of the First International Symposium on Soqatra Island: Present and Future* 1: 83-105. United Nations Publications, p. 219-232, New York.
- Schneider, W. & M.J. Parr 1998. *Orthetrum julia falsum* Longfield 1955, new to the dragonfly fauna of Yemen and the Arabian Peninsula (Anisoptera: Libellulidae). *International Journal of Odonatology* 1: 155-158.
- van der Weide, M.J.T. & V.J. Kalkman 2008. Some new records of dragonflies from Oman. *Agrion* 12: 52-54.
- Waterston, A.R. 1980. Insects of Saudi Arabia. Odonata. *Fauna of Saudi Arabia* 2: 57-70.
- *Waterston, A.R. 1984. A new genus and species of platycnemidid dragonfly from the Arabian peninsula (Zygoptera). *Odonatologica* 13: 139-146.
- Waterston, A.R. 1985. Insects of southern Arabia. Odonata from the Yemens and Saudi Arabia. *Fauna of Saudi Arabia* 6: 451-472.
- Waterston, A.R. & A.R. Pittaway 1991. The Odonata or dragonflies of Oman and neighbouring territories. *Journal of Oman Studies* 10 (for 1989): 131-168.
- Wilson, K. 2008: A brief trip to United Arab Emirates and northern Oman. *Agrion* 12: 56-57.

